Quantum Gravity as Theory of "Superfluidity"

B.M. Barbashov¹, V.N. Pervushin¹, A.F. Zakharov^{1,2}, and V.A. Zinchuk¹,

1. Joint Institute for Nuclear Research,

141980, Dubna, Russia.

2. National Astronomical Observatories of Chinese Academy of Sciences, Beijing 100012, China

Abstract

A version of the cosmological perturbation theory in general relativity (GR) is developed, where the cosmological scale factor is identified with spatial averaging of the metric determinant logarithm and the cosmic evolution acquires the pattern of a superfluid motion: the absence of "friction-type" interaction, the London-type wave function, and the Bogoliubov condensation of quantum universes. This identification keeps the number of variables of GR and leads to a new type of potential perturbations. A set of arguments is given in favor of that this "superfluid" version of GR is in agreement with the observational data.

> "Long Communication" at the XXVIII Spanish Relativity Meeting "A Century of Relativity Physics" Oviedo, Spain, 6-10 Sept. 2005 http://fisi24.ciencias.uniovi.es/ere05.html To be published in "AIP Conference Proceedings"

Contents

1.	Introduction	2
2.	"Superfluid" motion and potential perturbations	2
3.	London wave function & Bogoluibov condensation	5
4.	W-,Z- Factory versus the Planck epoch	5
5.	Conclusions	7

1. Introduction

Separation of the cosmological scale factor from metrics in general relativity (GR) is well known as the cosmological perturbation theory proposed by Lifshitz [1] and applied as a basic tool for analysis of modern observational data in astrophysics and cosmology [2]. However, the number of variables of the Lifshitz theory differs from GR. In the present paper, a new version of the cosmological perturbation theory [3] is discussed, where the logarithm of cosmological scale factor coincides with spatial averaging the metric determinant logarithm, so that the number of variables of GR is conserved. The conservation of the number of variables allows us to solve the energy constraint, fulfil the Dirac Hamiltonian reduction and quantization. This Quantum Gravity has some attributes of the theory of superfluidity of the type of Landau's "superfluid" dynamics [4], London's unique wave function [5], and Bogoliubov's squeezed condensate [6] and gives us a new possibility to explain the "CMBR primordial power spectrum" and other topical problems of modern cosmology.

2. "Superfluid" motion and potential perturbations

GR was given by Einstein and Hilbert 90 years ago with the "dynamic" action

$$S = \int d^4x \sqrt{-g} \left[-\frac{\varphi_0^2}{6} R(g) + \mathcal{L}_{(\mathrm{M})} \right], \tag{1}$$

where $\varphi_0^2 = \frac{3}{8\pi} M_{\text{Planck}}^2$ is the Newton constant, $\mathcal{L}_{(M)}$ is the matter Lagrangian, and an "interval" $g_{\mu\nu} dx^{\mu} dx^{\nu} \equiv \omega_{(\alpha)} \omega_{(\alpha)} = \omega_{(0)} \omega_{(0)} - \omega_{(1)} \omega_{(1)} - \omega_{(2)} \omega_{(2)} - \omega_{(3)} \omega_{(3)}$ presented here by the components of an orthogonal simplex of reference $\omega_{(\alpha)}$.

These simplex components in the reference frame of the Dirac - ADM Hamiltonian approach to GR [7] take the form

$$\omega_{(0)} = \psi^6 N_{\rm d} dx^0, \qquad \omega_{(b)} = \psi^2 \mathbf{e}_{(b)i} (dx^i + N^i dx^0); \qquad (2)$$

here $\mathbf{e}_{(a)i}$ are triads of the spatial metrics with det $|\mathbf{e}| = 1$, N_d is the Dirac lapse function, N^i is shift vector, and ψ is a determinant of the spatial metric. All these components were considered by Dirac as the potentials except of two transverse triads distinguished by the constraints $\partial_j \mathbf{e}_{(a)}^j = 0$ and the zero determinant momentum $p_{\psi} = 0$. Latter contradicts to Friedmann-type evolution in the homogeneous approximation of GR, where the determinant component is identified with the cosmological scale factor ($\psi^2 \simeq a(x^0)$) and its momentum coincides with the Hubble parameter. The question appears about the status of the Hubble parameter in the Dirac Hamiltonian approach [7].

To answer this question we consider the cosmological perturbation theory [1, 2] $g_{\mu\nu} = a^2(x^0)\tilde{g}_{\mu\nu}$ in terms of the Dirac – ADM variables, where one can get the new lapse function $\tilde{N}_d = [\sqrt{-\tilde{g}} \ \tilde{g}^{00}]^{-1} = a^2 N_d$ and spatial determinant $\tilde{\psi} = (\sqrt{a})^{-1} \psi$.

After the separation of the spatial metric determinant in the curvature $\sqrt{-g}R(g) = a^2\sqrt{-\tilde{g}}R(\tilde{g}) - 6a\partial_0\left[\widetilde{N}_d^{-1}\partial_0 a\right]$ the determinant part of the Lagrangian takes the form

$$L = -\varphi_0^2 \int d^3 x \widetilde{N}_d \left[4a^2 \left(\overline{v_\psi} \right)^2 + \left(\frac{\partial_0 a}{\widetilde{N}_d} \right)^2 \right] - \left[2\varphi_0^2 \partial_0 a^2 \int d^3 x \overline{v_\psi} \right] + \dots, \tag{3}$$

where $\overline{v_{\psi}} = \left[(\partial_0 - N^l \partial_l) \log \widetilde{\psi} - \frac{1}{6} \partial_l N^l \right] / \widetilde{N}_d$ is a local velocity. One can see that this Lagrangian contains the velocity-velocity interaction distinguished by the box in Eq. (3). This interaction

mixes the canonical momenta

$$P_a \equiv \frac{\partial L}{\partial(\partial_0 a)} = -\varphi_0^2 \int d^3 x \left[\boxed{4a \ \overline{v_\psi}} + 2 \frac{\partial_0 a}{\widetilde{N}_d} \right], \tag{4}$$

$$\overline{p_{\psi}} = \frac{\partial \mathcal{L}}{\partial(\partial_0 \log \tilde{\psi})} = -2\varphi_0^2 a \left[4a\overline{v_{\psi}} + \left[2 \frac{\partial_0 a}{\tilde{N}_d} \right] \right], \tag{5}$$

so that the integral of the local momentum $\overline{p_{\psi}}$ coincides with the one of cosmic motion

$$\int d^3x \overline{p_{\psi}} = -2\varphi_0^2 a \int d^3x \left[4a\overline{v_{\psi}} + \left[2 \frac{\partial_0 a}{\widetilde{N_d}} \right] \right] \equiv \boxed{2aP_a},\tag{6}$$

and the Hamiltonian approach is failure. The reason of that is the increase of the number of variables after the separation $a(x^0)$.

In order to keep the number of variables, we identify $\log \sqrt{a}$ with the spatial volume "averaging" of $\log \psi$: $\log \sqrt{a} = \langle \log \psi \rangle \equiv \int d^3x \log \psi / V_0$, where $V_0 = \int d^3x < \infty$ is a finite volume. In this case, the new determinant variable $\tilde{\psi}$ and its velocity satisfy the identities

$$\int d^3x \log \widetilde{\psi} = \int d^3x \left[\log \psi - \langle \log \psi \rangle\right] \equiv 0, \qquad \int d^3x \overline{v_\psi} \equiv 0. \tag{7}$$

This means that all the box terms in Eqs (3)- (6) disappear together with the "friction-type" velocity-velocity interaction. The momenta (4) and (5) are completely separated

$$P_a \equiv -2V_0\varphi_0^2 \partial_0 a \langle (\widetilde{N}_d)^{-1} \rangle = -2V_0\varphi_0^2 \frac{da}{d\zeta} \equiv -2V_0\varphi_0^2 a', \tag{8}$$

$$\overline{p_{\psi}} = \frac{\partial \mathcal{L}}{\partial(\partial_0 \log \widetilde{\psi})} = -8\varphi_0^2 a^2 \overline{v_{\psi}}, \qquad (9)$$

here the averaging $\langle (\tilde{N}_d)^{-1} \rangle$ determines the time interval $d\zeta = \langle (\tilde{N}_d)^{-1} \rangle^{-1} dx^0$ that is invariant with respect to reparametrizations of the coordinate evolution parameter x^0 . In this case, the scale transformation $g_{\mu\nu} = a^2(x^0)\tilde{g}_{\mu\nu}$ converts the GR action (1) into the "friction-free" one

$$S[\varphi_0] = \widetilde{S}[\varphi] - \int dx^0 (\partial_0 \varphi)^2 \int \frac{d^3 x}{\widetilde{N}_d},$$
(10)

where $\widetilde{S}[\varphi]$ is the action (1) in terms of metrics \widetilde{g} and the running scale $\varphi(x^0) = \varphi_0 a(x^0)$ of all masses including the Planck one φ_0 . The energy constraint $\delta S[\varphi_0]/\delta \widetilde{N}_d = 0$ takes algebraic form [9] with respect to the invariant local lapse function $N_{\text{inv}} \equiv \widetilde{N}_d \langle (\widetilde{N}_d)^{-1} \rangle$

$$-\frac{\delta \widetilde{S}[\varphi]}{\delta \widetilde{N}_d} \equiv \widetilde{T}_0^0 = \left[\frac{\partial_0 \varphi}{\widetilde{N}_d}\right]^2 = \frac{1}{N_{\rm inv}^2} \left[\frac{d\varphi}{d\zeta}\right]^2 = \frac{\varphi'^2}{N_{\rm inv}^2},\tag{11}$$

where \widetilde{T}_0^0 is the local energy density. This equation has the resolution in both the local sector

$$N_{\rm inv} \equiv \widetilde{N}_d \langle (\widetilde{N}_d)^{-1} \rangle = \left\langle \sqrt{\widetilde{T}_0^0} \right\rangle \left(\sqrt{\widetilde{T}_0^0} \right)^{-1} \tag{12}$$

and the global one

$$\zeta(\varphi_0|\varphi) \equiv \int dx^0 \langle (\widetilde{N}_d)^{-1} \rangle^{-1} = \pm \int_{\varphi}^{\varphi_0} d\widetilde{\varphi} \left\langle \sqrt{\widetilde{T}_0^0(\widetilde{\varphi})} \right\rangle^{-1}$$
(13)

where the invariant time ζ is connected with the scale factor by the Hubble-like law.

The explicit dependence of \widetilde{T}_0^0 on $\widetilde{\psi}$ was given by Lichnerowicz [10]

$$\widetilde{T}_{0}^{0} = \widetilde{\psi}^{7} \triangle \widetilde{\psi} + \sum_{I} \widetilde{\psi}^{I} a^{\frac{I}{2} - 2} \tau_{I}, \qquad \tau_{I} \equiv \langle \tau_{I} \rangle + \overline{\tau_{I}}, \qquad (14)$$

where $\hat{\Delta}\tilde{\psi} \equiv \frac{4\varphi^2}{3}\partial_{(b)}\partial_{(b)}\tilde{\psi}$ is the Laplace operator and τ_I is partial energy density marked by the index I running a set of values I = 0 (stiff), 4 (radiation), 6 (mass), 8 (curvature), 12 (Aterm) in correspondence with a type of matter field contributions. The negative contribution $-(16/\varphi^2)\overline{p_{\psi}}^2$ of the spatial determinant momentum in the energy density $\tau_{I=0}$ can be removed by the Dirac constraint [7] of the zero velocity of the spatial volume element

$$\overline{p_{\psi}} = -8\varphi^2 \frac{1}{\widetilde{N}_d} \left[(\partial_0 - N^l \partial_l) \ln \widetilde{\psi} - \frac{1}{6} \partial_l N^l \right] = -8\varphi^2 \frac{\partial_0 \widetilde{\psi}^6 - \partial_l [\widetilde{\psi}^6 N^l]}{\widetilde{\psi}^6 \widetilde{N}_d} = 0.$$
(15)

In the class of functions $\overline{F} = F - \langle F \rangle$, the classical equation $\delta S / \delta \log \widetilde{\psi} = 0$ takes the form

$$\widetilde{N_d}\widetilde{\psi}\frac{\partial\widetilde{T}_0^0}{\partial\widetilde{\psi}} + \widetilde{\psi}\triangle\left[\frac{\partial\widetilde{T}_0^0}{\partial\triangle\widetilde{\psi}}\widetilde{N_d}\right] = 0 \quad \to \quad 7\overline{N_{\rm inv}\widetilde{\psi}^7\hat{\bigtriangleup}\widetilde{\psi}} + \overline{\widetilde{\psi}\widehat{\bigtriangleup}[N_{\rm inv}\widetilde{\psi}^7]} + \sum_I I\overline{\widetilde{\psi}^I a^{\frac{I}{2}-2}\tau_I} = 0$$

and gives both the determinant $\tilde{\psi} = 1 + \overline{\mu}$ and the lapse function (12) $N_{\text{inv}}\tilde{\psi}^7 = 1 - \overline{\nu}$.

The first order of this equation determines $\overline{\mu}$ and $\overline{\nu}$ in the form of a sum [3]

$$\overline{\mu} = \frac{1}{2} \int d^3y \left[D_{(+)}(x,y) \overline{T}_{(+)}^{(\mu)}(y) + D_{(-)}(x,y) \overline{T}_{(-)}^{(\mu)}(y) \right],$$
(16)

$$\overline{\nu} = \frac{1}{2} \int d^3 y \left[D_{(+)}(x,y) \overline{T}_{(+)}^{(\nu)}(y) + D_{(-)}(x,y) \overline{T}_{(-)}^{(\nu)}(y) \right], \tag{17}$$

where

$$\beta = \sqrt{1 + [\langle \tau_{(2)} \rangle - 14 \langle \tau_{(1)} \rangle] / (98 \langle \tau_{(0)} \rangle)},$$

$$\overline{T}_{(\pm)}^{(\mu)} = [1 \pm 49\beta]\overline{\tau_{(0)}} \mp 7\beta\overline{\tau_{(1)}}, \qquad \overline{T}_{(\pm)}^{(\nu)} = [7 \pm (14\beta)^{-1}]\overline{\tau_{(0)}} - \overline{\tau_{(1)}}$$
(18)

are the local currents, $D_{(\pm)}(x,y)$ are the Green functions satisfying the equations

$$[\pm \hat{m}_{(\pm)}^2 - \hat{\Delta}] D_{(\pm)}(x, y) = \delta^3(x - y), \tag{19}$$

where $\hat{m}_{(\pm)}^2 = 14(\beta \pm 1)\langle \tau_{(0)} \rangle \mp \langle \tau_{(1)} \rangle$, $\tau_{(n)} = \sum_I I^n a^{\frac{I}{2}-2} \tau_I$. In the case of point mass distribution in a finite volume V_0 with the zero pressure and the density $\overline{\tau_{(0)}}(x) = \overline{\tau_{(1)}}(x)/6 \equiv M \left[\delta^3(x-y) - 1/V_0 \right]$, solutions (16), (17) take a very interesting form

$$\widetilde{\psi} = 1 + \overline{\mu}(x) = 1 + \frac{r_g}{4r} \left[\gamma_1 e^{-m_{(+)}(z)r} + (1 - \gamma_1) \cos m_{(-)}(z)r \right],$$
(20)

$$N_{\rm inv}\widetilde{\psi}^7 = 1 - \overline{\nu}(x) = 1 - \frac{r_g}{4r} \left[(1 - \gamma_2)e^{-m_{(+)}(z)r} + \gamma_2 \cos m_{(-)}(z)r \right], \tag{21}$$

where $\gamma_1 = \frac{1+7\beta}{2}$, $\gamma_2 = \frac{14\beta - 1}{28\beta}$, $r_g = \frac{3M}{4\pi\varphi^2}$, r = |x - y|. The zero volume velocity (15) gives the diffeo-invariant shift of the coordinate origin

$$\langle (\widetilde{N}_d)^{-1} \rangle N^i = \left(\frac{x^i}{r}\right) \left(\frac{\partial_{\zeta} V}{\partial_r V}\right), \qquad V(\zeta, r) = \int^r d\widetilde{r} \ \widetilde{r}^2 \widetilde{\psi}^6(\zeta, \widetilde{r}). \tag{22}$$

In the infinite volume limit $\langle \tau_{(n)} \rangle = 0$, a = 1 solutions (20) and (21) coincide with the isotropic version of the Schwarzschild solutions: $\tilde{\psi} = 1 + \frac{r_g}{4r}$, $N_{\text{inv}}\tilde{\psi}^7 = 1 - \frac{r_g}{4r}$, $N^k = 0$.

The main differences of the superfluid-type version of GR from the Lifshitz version [1, 2] are the potential perturbations of the scalar components N_{inv} , $\tilde{\psi}$ instead of the kinetic ones and the nonzero shift vector $N^k \neq 0$ determined by Eq. (15). Recall that just the kinetic perturbations are responsible for the "primordial power spectrum" in the inflationary model [2]. The problem appears to describe CMBR by the potential perturbations given by Eqs. (16) – (22).

3. London wave function & Bogoluibov condensation

The main consequence of the separation of the cosmological scale factor is the London-type globalization of the energy constraint (12). It fixes only the scale momentum P_{φ}

$$P_{\varphi}^2 \equiv [P_a/\varphi_0]^2 = E_{\varphi}^2 \equiv \left[2\int d^3x \sqrt{\widetilde{T}_0^0}\,\right]^2,\tag{23}$$

in contrast to the energy constraint in GR without the cosmological scale factor. The "reduced" action can be obtained as values of the Hamiltonian action for the energy constraint $P_{\varphi}^2 = E_{\varphi}^2$ [9]:

$$S[\varphi_I|\varphi_0]|_{P_{\varphi}=\pm E_{\varphi}} = \int_{\varphi_I}^{\varphi_0} d\widetilde{\varphi} \left\{ \int d^3x \left[\sum_F P_F \partial_{\varphi} F + C \mp 2\sqrt{\widetilde{T}_0^0(\widetilde{\varphi})} \right] \right\},\tag{24}$$

where C is the sum of all Dirac constraints [7, 9]. Momenta $P_{\varphi\pm} = \pm E_{\varphi}$ become the generator of evolution of all variables with respect to the evolution parameter φ [9] forward and backward, respectively. The negative energy problem can be solved by the primary quantization of the energy constraint $[P_{\varphi}^2 - E_{\varphi}^2]\Psi_u = 0$ and the secondary quantization $\Psi_u = (1/\sqrt{2E_{\varphi}})[A^+ + A^-]$ by the Bogoliubov transformation $A^+ = \alpha B^+ + \beta^* B^-$, in order to diagonalize the equations of motion by the condensation of "universes" $< 0|\frac{i}{2}[A^+A^+ - A^-A^-]|0 >= R(\varphi)$ and describe cosmological creation of a "number" of universes $< 0|A^+A^-|0 >= N(\varphi)$ from the stable Bogoliubov vacuum $B^-|0 >= 0$. Vacuum postulate $B^-|0 >= 0$ leads to an arrow of the invariant time $\zeta \geq 0$ (13) and its absolute point of reference $\zeta = 0$ at the moment of creation $\varphi = \varphi_I$ [3, 9]; whereas the Planck value of the running mass scale $\varphi_0 = \varphi(\zeta = \zeta_0)$ belongs to the present day moment ζ_0 . The reduced action (24) shows us that the initial data at the beginning $\varphi = \varphi_I$ are independent of the present-day ones at $\varphi = \varphi_0$, therefore the proposal about an existence of the Planck epoch $\varphi = \varphi_0$ at the beginning [2] looks very doubtful.

4. W-,Z- Factory versus the Planck epoch

The low-energy expansion of the "reduced action" (24) over the field density $T_{\rm sm}$

$$2d\varphi\sqrt{\widetilde{T}_0^0} = 2d\varphi\sqrt{\rho_0(\varphi) + T_{\rm sm}} = d\varphi\left[2\sqrt{\rho_0(\varphi)} + T_{\rm sm}/\sqrt{\rho_0(\varphi)}\right] + \dots$$

gives the sum: $S^{(+)}|_{\text{constraint}} = S^{(+)}_{\text{cosmic}} + S^{(+)}_{\text{field}} + \dots$, where $S^{(+)}_{\text{cosmic}}[\varphi_I|\varphi_0] = -2V_0 \int_{\varphi_I}^{\varphi_0} d\varphi \sqrt{\rho_0(\varphi)}$ is the reduced cosmological action and

$$S_{\text{field}}^{(+)} = \int_{\eta_I}^{\eta_0} d\eta \int d^3x \left[\sum_F P_F \partial_\eta F - T_{\text{sm}} \right]$$
(25)

is the standard field action in terms of the conformal time: $d\zeta = d\eta = d\varphi/\sqrt{\rho_0(\varphi)}$ in the conformal flat space-time with running masses $m(\eta) = a(\eta)m_0$.

This low-energy expansion identifies the "conformal quantities" with the observable ones including the conformal time $d\eta$, instead of $dt = a(\eta)d\eta$, the coordinate distance r, instead of Friedmann one $R = a(\eta)r$, and the conformal temperature $T_c = Ta(\eta)$, instead of the standard one T. In this case, the cosmological redshift of the spectral lines

$$\frac{E_{\text{emission}}}{E_0} = \frac{m_{\text{atom}}(\eta_0 - r)}{m_{\text{atom}}(\eta_0)} \equiv \frac{\varphi(\eta_0 - r)}{\varphi_0} = a(\eta_0 - r) = \frac{1}{1 + z}$$

is explained by the running masses $m = a(\eta)m_0$ in action (25) [11, 12, 13].

The conformal observable distance r loses the factor a, in comparison with the nonconformal one R = ar. In the units of "conformal quantities" the Supernova data [15, 16] are consistent with the dominance of the stiff (rigid) state, $\Omega_{\text{Rigid}} \simeq 0.85 \pm 0.15$, $\Omega_{\text{Matter}} = 0.15 \pm 0.15$ [12, 13]. If $\Omega_{\text{Rigid}} = 1$, we have the square root dependence of the scale factor on conformal time $a(\eta) = \sqrt{1 + 2H_0(\eta - \eta_0)}$. Just this time dependence of the scale factor on the measurable time (here – conformal one) is used for description of the primordial nucleosynthesis [17]. This stiff state is formed by a free scalar field when $E_{\varphi} = 2V_0\sqrt{\rho_0} = Q/\varphi$ and $Q/2V_0 = H_0\varphi_0^2 = H_I\varphi_I^2$ is an integral of motion.

These initial data φ_I and H_I can be determined by the parameters of matter cosmologically created from the vacuum at the beginning of a universe $\eta \simeq 0$.

The Standard Model (SM) density $T_{\rm sm}$ in action (25) shows us that W-, Z- vector bosons have maximal probability of this cosmological creation due to their mass singularity [14]. One can introduce the notion of a particle in a universe if the Compton length of a particle defined by its inverse mass $M_{\rm I}^{-1} = (a_{\rm I}M_{\rm W})^{-1}$ is less than the universe horizon defined by the inverse Hubble parameter $H_{\rm I}^{-1} = a_{\rm I}^2(H_0)^{-1}$ in the stiff state. Equating these quantities $M_{\rm I} = H_{\rm I}$ one can estimate the initial data of the scale factor $a_{\rm I}^2 = (H_0/M_{\rm W})^{2/3} = 10^{-29}$ and the primordial Hubble parameter $H_{\rm I} = 10^{29}H_0 \sim 1 \text{ mm}^{-1} \sim 3K$. Just at this moment there is an effect of intensive cosmological creation of the vector bosons described in [14]; in particular, the distribution functions of the longitudinal vector bosons demonstrate us a large contribution of relativistic momenta. Their conformal (i.e. observable) temperature T_c (appearing as a consequence of collision and scattering of these bosons) can be estimated from the equation in the kinetic theory for the time of establishment of this temperature $\eta_{\rm relaxation}^{-1} \sim n(T_c) \times \sigma \sim H$, where $n(T_c) \sim T_c^3$ and $\sigma \sim 1/M^2$ is the cross-section. This kinetic equation and values of the initial data $M_{\rm I} = H_{\rm I}$ give the temperature of relativistic bosons

$$T_c \sim (M_{\rm I}^2 H_{\rm I})^{1/3} = (M_0^2 H_0)^{1/3} \sim 3K$$
 (26)

as a conserved number of cosmic evolution compatible with the Supernova data [12, 15, 16]. We can see that this value is surprisingly close to the observed temperature of the CMB radiation $T_c = T_{\text{CMB}} = 2.73$ K.

The primordial mesons before their decays polarize the Dirac fermion vacuum (as the origin of axial anomaly) and give the baryon asymmetry frozen by the CP-violation. The value of the baryon-antibaryon asymmetry of the universe following from this axial anomaly was estimated in [14] in terms of the coupling constant of the superweak-interaction

$$n_b/n_\gamma \sim X_{CP} = 10^{-9}.$$
 (27)

The boson life-times $\tau_W = 2H_I \eta_W \simeq (2/\alpha_W)^{2/3} \simeq 16$, $\tau_Z \sim 2^{2/3} \tau_W \sim 25$ determine the presentday visible baryon density

$$\Omega_b \sim \alpha_W = \alpha_{QED} / \sin^2 \theta_W \sim 0.03. \tag{28}$$

All these results (26) – (28) testify to that all visible matter can be a product of decays of primordial bosons, and the observational data on CMBR can reflect parameters of the primordial bosons, but not the matter at the time of recombination. In particular, the length of the semicircle on the surface of the last emission of photons at the life-time of W-bosons in terms of the length of an emitter (i.e. $M_W^{-1}(\eta_L) = (\alpha_W/2)^{1/3}(T_c)^{-1}$) is $\pi \cdot 2/\alpha_W$. It is close to $l_{min} \sim 210$ of CMBR, whereas $(\Delta T/T)$ is proportional to the inverse number of emitters $(\alpha_W)^3 \sim 10^{-5}$.

The temperature history of the expanding universe copied in the "conformal quantities" looks like the history of evolution of masses of elementary particles in the cold universe with the conformal temperature $T_c = a(\eta)T$ of the cosmic microwave background. In the superfluid version of cosmology [12], the dominance of stiff state $\Omega_{\text{Stiff}} \sim 1$ determines the parameter of spatial oscillations $\hat{m}^2_{(-)} = \frac{6}{7}H_0^2[\Omega_{\text{R}}(z+1)^2 + \frac{9}{2}\Omega_{\text{Mass}}(z+1)]$. The values of red shift in the recombination epoch $z_r \sim 1100$ and the clusterization parameter $r_{\text{clustering}} = \frac{\pi}{\hat{m}_{(-)}} \sim \frac{\pi}{H_0\Omega_R^{1/2}(1+z_r)} \sim 130 \,\text{Mpc}$ re-

cently discovered in studies of a large scale periodicity in redshift distribution [18] lead to a reasonable value of the radiation-type density $10^{-4} < \Omega_R \sim 3 \cdot 10^{-3} < 5 \cdot 10^{-2}$ at the time of this epoch.

5. Conclusions

The conservation of the number of variables of GR after the separation of the cosmological scale factor from all fields leads to the Hamiltonian version of GR where the cosmic evolution acquires the pattern of a superfluid motion without the friction-type interaction. In this version the scale factor is an evolution parameter in the "field space of events" $[\varphi|\tilde{F}^{(n)}]$, and its canonical momentum (i.e. the Hubble "parameter") plays the role of the generator of evolution of the fields $\tilde{F}^{(n)}$. The values of the scale factor momentum for solutions of the equations of motion can be called the "reduced energies".

The solution of the problem of the "negative reduce energy" by the primary quantization and the secondary one (on the analogy of the pathway passed by QFT in the 20th century) reveals in GR other attributes of the theory of superfluid quantum liquid: London-type WDW wave function and Bogoliubov-type condensate of quantum universes. The postulate of the quantum Bogoliubov vacuum as the state with the minimal "energy" leads to the absolute beginning of geometric time.

The Hamiltonian approach leads to potential perturbations of the scalar metric components in contrast to the standard cosmological perturbation theory [1] keeping only the kinetic perturbations which are responsible for the "primordial power spectrum" in the inflationary model [2]. The Quantum Gravity considered as the theory of superfluidity gives us a possibility to explain this "spectrum" and other topical problems of cosmology by the cosmological creation of the primordial W-, Z- bosons from vacuum, when their Compton length coincides with the universe horizon.

The correspondence principle as the low-energy expansion of the reduced action identifies the conformal quantities with the "measurable" ones, and the uncertainty principle establishes the point of the beginning of the cosmological creation of the primordial W-, Z- bosons from vacuum due to their mass singularity at the moment $a_I^2 \simeq 10^{-29}$, $H_I^{-1} \simeq 1$ mm. In this case, the equations describing the longitudinal vector bosons in SM are close to the equations of the inflationary model used for description of the "power primordial spectrum" of the CMB radiation [2]. We listed the set of theoretical and observational arguments in favor of that the CMB radiation can be a final product of primordial vector W-, Z- bosons cosmologically created from the vacuum.

Acknowledgement

The authors are grateful to A.A. Gusev, A.V. Efremov, E.A. Kuraev, V.V. Nesterenko, V.B. Priezzhev, and S.I. Vinitsky for interesting and critical discussions. AFZ is grateful to the National Natural Science Foundation of China (NNSFC) (Grant # 10233050) for a partial financial support.

References

- E.M. Lifshitz, ZhETF, 16, 587-602 (1946); J.M. Bardeen, Phys. Rev. D, 22, 1882-1905 (1980).
- [2] V. F. Mukhanov, H. A. Feldman, and R. H. Brandenberger, Phys. Rep. 215, 206–333 (1992).
- [3] V.N. Pervushin and V.A. Zinchuk, gr-qc/0504123;
 B.M. Barbashov, V.N. Pervushin, A.F. Zakharov, and V.A. Zinchuk, gr-qc/0509006.
- [4] L.D. Landau, *ZhETF*, **11**, 592–614 (1941).
- [5] F. London, *Nature*, **141**, 643–645 (1938).
- [6] N.N. Bogoliubov, J. Phys., 11, 23-32 (1947).
- [7] P.A.M. Dirac, Proc. Roy. Soc., A 246, 333-344 (1958); P.A.M. Dirac, Phys. Rev., 114, 924-930 (1959); R. Arnovitt, S. Deser, and C.W. Misner, Phys. Rev., 117, 1595-1602 (1960).
- [8] A.L. Zelmanov, Dokl. AN USSR, 107, 815–818 (1956); Dokl. AN USSR, 209, 822–825 (1973).
- [9] M. Pawlowski and V.N. Pervushin, Int. J. Mod. Phys., 16, 1715–1742 (2001); [hep-th/0006116];
 B.M. Barbashov, V.N. Pervushin, and D.V. Proskurin, Theor. Math. Phys., 132, 1045–1058 (2002).
- [10] A. Lichnerowicz, Journ. Math. Pure and Appl., 23, 37 47 (1944).
- [11] J.V. Narlikar, Introduction to Cosmology, Jones and Bartlett, Boston, 1983.
- [12] D. Behnke, D.B. Blaschke, V.N. Pervushin, and D.V. Proskurin, *Phys. Lett. B*, **530**, 20–26 (2002); [gr-qc/0102039].
- [13] D. Behnke, *PhD Thesis*, Rostock Report MPG-VT-UR 248/04 (2004)
- [14] D.B. Blaschke et al., Physics of Atomic Nuclei, 67, 1050 1062 (2004); [hep-ph/0504225].
- [15] A.G. Riess et al., Astron. J., 116, 1009–1038 (1998); S. Perlmutter et al., Astrophys. J., 517, 565–586 (1999); A.G. Riess et al., Astrophys. J., 607, 665–687 (2004).
- [16] A.G. Riess et al., Astrophys. J., 560, 49–71 (2001).
- [17] S. Weinberg, First Three Minutes. A Modern View of the Origin of the Universe, Basic Books, Inc., Publishers, New-York, 1977.
- [18] W.J. Cocke and W.G. Tifft, Astrophys. J., 368, 383–389 (1991).